



Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)
Please Send a Copy of The Patient's Up-to-Date Clinical Notes

PATIENT INFORMATION (Complete or Fax Existing Chart)				PRESCRIBER INFORMATION			
Name:				State License:			
INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back)							
Primary Insurance: Plan #: Group #: RX Card (PBM): BIN:				Plan #: Group #: RX Card (PBM):			
CLINICAL INFORMATION							
Primary ICD-10 (Please Specify Diagnosis): Secondary ICD-10 (Please Specify Diagnosis): Is the patient on iron, folate and/or vitamin B12 therapy?							
ARANESP® ORDERS							
Prescription type: New start Restart Continued therapy Total Doses Received: Date of Last Injection/Infusion:							
Medication ☐ Aranesp® (darbepoetin alfa) Single Dose Vials ☐ Aranesp® (darbepoetin alfa) Single Dose Prefilled Syringe		☐ 150 mcg ☐	340 mcg				
Special Instructions:							
Pre-Medication		Dose/Strength		Directions			
☐ Acetaminophen ☐ Diphenhydramine	☐ 500mg ☐ 25mg IV/PO ☐ 50mg IV/PO			☐ Take 1-2 tablets PO prior to infusion or post-infusion as directed ☐ Take 1 tablet PO prior to infusion or as directed OR ☐ Inject contents of 1 vial IV prior to infusion or as directed			
☐ Methylprednisolone	☐ 40mg ☐ 125mg			☐ Inject contents of 1 vial IV prior to infusion or as directed			
INFUSION REACTION ORDERS							
Mild reaction protocol: ☑ Diphenhydramine 25mg IV, one time, for pruritus. If symptoms worsen, see orders for moderate to severe reactions. Moderate reaction protocol:							

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.



medicine as prescribed in this referral.

ARANESP®

Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)
Please Send a Copy of The Patient's Up-to-Date Clinical Notes

Date: __

☑ Acetaminophen 650mg PO, one time, for pyrexia or rigors							
☑ Diphenhydramine 50mg IV, one time, for pruritus or urticaria							
☑ Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms							
If symptoms worsen, see interventions for	or severe reactions						
Severe reaction protocol: (Call 911 if init	tiated):						
☑ Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%)							
☐ Diphenhydramine 50mg IV,one time, for respiratory symptoms, edema, or anaphylaxis							
☑ Methylprednisolone 125mg IV, one time, for respiratory symptoms, edema, or anaphylaxis							
☑ Sodium Chloride 0.9% 500mL IV over 30-60 min, one time, for cardiovascular symptoms							
☑ Epinephrine 0.3mg/0.3mL IM into mis-anterolateral aspect of thigh of anaphylaxis, may repeat x1 in 5-15 minutes if symptoms are not resolved or							
worsen							
FLUSHING & LOCKING ORDERS							
Flushing Protocol (>66lbs/33kg)							
PIV and Midline:		Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter:					
☑ 0.9% Sodium Chloride 2-5mL IV flush	before and after each	☑ 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mL IV flush					
infusion		after infusion/lab draw					
Locking Protocol (>66lbs/33kg)							
PIV and Midline:	PICC:		Implanted Port, Tunneled Catheter, and Non-tunneled Catheter:				
☐ Heparin Sodium 10 units/mL 1mL	⊠ Heparin Sodium 10 un	nits/mL 3mL IV	☐ Heparin Sodium 100 units/mL 3-5mL IV final flush post normal				
IV final flush post normal saline flush	final flush post normal sa	aline flush	saline flush				
** May substitute Dextrose 5% in Water, or alternative, for 0.9& Sodium Chloride, when indicated due to incompatibility with medications bring infused							
SIGNATURE							

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

We hereby authorize Talis Healthcare LLC to provide all supplies and additional services (nursing/patient training) required to provide and deliver the

Prescriber Signature

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.